



LARC Quick Coding Guide Supplement



LARC ICD-10 Coding Reference

Basic Implant Codes

ICD-10 Codes – Implant		CPT Procedure Codes	HCPCSII/JCode
Z30.017	Encounter for initial prescription and insertion of subdermal contraceptive, even when device is placed at a separate encounter	11981 - insertion	J7307
Z30.46	Encounter for surveillance of subdermal contraceptive (includes removal, checking, reinsertion)	11982 – removal 11983 – removal + reinsertion	J7307
Z32.02	Pregnancy test/exam – negative	81025 (Urine pregnancy test)	Don't forget to bill for the point-of-care office pregnancy test (when conducted)

Basic IUD Codes

ICD-10 Codes - IUDs		CPT Procedure Code	HCPCSII/JCode
Z30.014	Encounter for initial prescription of IUD (not coded with the actual insertion)		
Z30.430	Encounter for insertion of IUD	58300	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Z30.431	Follow-up for patient with IUD or Routine checking for IUD		
Z30.432	Encounter for removal of IUD	58301	
Z30.433	Encounter for removal + reinsertion of IUD	58301 AND 58300-51* OR 58300-59* (Check with payer for expected modifier.) Append modifier -51 or -59 to the lesser paying service.	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Z32.02	Pregnancy test/exam – negative	81025 (Urine pregnancy test)	Don't forget to bill for the point-of-care office pregnancy test (when conducted)

LARC Quick Coding Guide Supplement

Coding Complex Cases (IUDs) with ICD-10

Clinical scenario	ICD-10 Codes		CPT Procedure Code	HCPCSII/JCode
Failed insertion/ discontinued procedure	Z30.430 AND <Co-occurring complication which caused failed/ discontinued procedure>	Encounter for insertion of IUD *Document reason for failed/stopped procedure with appropriate ICD-10 codes.	58300-52* or -53* *NOTE: Use modifier -52 (Failed Procedure) to denote that you attempted insertion but the procedure was incomplete due to anatomical factors (eg. Stenosis) or -53 (Discontinued Procedure) to indicate that you had to stop because of concerns for patient well-being (eg. vaso-vagal, severe pain).	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Perforation (during insertion procedure)	Z30.430 T83.39XA	Encounter for insertion of IUD Perforation of uterus by IUD (non-traumatic)	58300-53 *Use modifier -53 if procedure was discontinued due to perforation.	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Difficult insertion with ultrasound guidance	Z30.430 AND <Co-occurring complication justifying ultrasound>	Encounter for insertion of IUD	58300-22 *Document the reason for additional work. 76998 (Ultrasonic guidance, intraoperative) *Document the justification for ultrasonic guidance (eg. patient in severe pain).	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301

LARC Quick Coding Guide Supplement

Clinical scenario	ICD-10 Codes		CPT Procedure Code	HCPCSII/JCode
Difficult insertion with ultrasound used to confirm the location of the IUD	Z30.430 <Co-occurring complication justifying ultrasound>	Encounter for insertion of IUD *Document complication with appropriate ICD-10 codes.	58300-22 *Document the reason for additional work. 76857 Ultrasound, pelvic [nonobstetric], real time with image documentation; limited or follow-up -or - 76830 Ultrasound, transvaginal *NOTE: It is not routine practice to use ultrasound to confirm placement. You must document justification for ultrasonography (eg. Uterine perforation, severe pain).	Kyleena = J7296 Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Missing strings, with ultrasound to locate	T83.32XA AND either Z30.431 OR Z30.432	Displacement of IUD - Missing strings, initial encounter	76857 Ultrasound, pelvic, limited or follow-up - OR - 76830 Ultrasound, transvaginal	
	Z30.431	Follow-up for patient with IUD (if patient wants to keep IUD)		
	Z30.432	Encounter for IUD removal (if patient desires removal)	58301 or 58301-22* *NOTE: You may append modifier -22 if the removal was complicated. Supporting documentation may be requested by a payer. If the string is easily located in the canal, -22 modifier should <i>not</i> be added. It should be appended only if it is a very difficult extraction and is separately documented with the claim.	

[illegible]

LARC Quick Coding Guide Supplement

Common LARC Modifiers

Modifier	Definition	Possible Clinical Scenarios	Documentation
-22	Increased procedural services	<ul style="list-style-type: none"> Complex or difficult insertion Unsuccessful insertion, followed by successful insertion during the same surgical session 	<ul style="list-style-type: none"> Total time of the procedure as compared with typical duration Reason for the additional work required Include diagnoses with appropriate ICD-10 codes or simple descriptive diagnoses that explain the reasons for the added difficulty
-25	Significant, separately identifiable E/M service	<ul style="list-style-type: none"> Patient came in for general contraceptive counseling, ends up choosing IUD or implant, and it is inserted that day 	<ul style="list-style-type: none"> Document time spent counseling
-51*	Multiple procedures performed on the same day, during the same session	<ul style="list-style-type: none"> Removal of IUD and insertion of new IUD on the same day Removal of implant and insertion of new implant on the same day 	<ul style="list-style-type: none"> Document should support the reasons for removal and reinsertion on the same day (eg. IUD expired, desired to continue with same method)
-52	Failed procedure	<ul style="list-style-type: none"> Provider couldn't complete procedure for anatomic reasons (eg. stenosis) 	<ul style="list-style-type: none"> Document reasons for procedure failure (eg. N88.2 Stricture/ stenosis of cervix)
-53	Discontinued procedure	<ul style="list-style-type: none"> Provider couldn't complete procedure due to concerns for patient well-being Severe pain Vasovagal Patient changed mind during procedure 	<ul style="list-style-type: none"> How much work was actually performed Document reason procedure terminated (eg. R55 Syncope/ vasovagal)
-59*	Distinct procedural service	<ul style="list-style-type: none"> Removal of IUD and insertion of new IUD on the same day Removal of implant and insertion of new implant on the same day 	<ul style="list-style-type: none"> Document should support the reasons for removal and reinsertion on the same day (eg. IUD expired, desired to continue with same method)
-76	Repeat procedure	<ul style="list-style-type: none"> Successful insertion but the IUD is expelled, followed by repeat insertion 	<ul style="list-style-type: none"> Document reason for repeat procedure (eg. IUD was expelled)

* When choosing between modifiers -51 and -59, payer policy may be the determining factor. Some payers will not pay for multiple procedures using modifier -51. **Check with payer.**